

# Local uniqueness for the Dirichlet-to-Neumann map via the two-plane transform

Allan Greenleaf\*

Department of Mathematics  
University of Rochester  
Rochester, NY 14627

Gunther Uhlmann†

Department of Mathematics  
University of Washington  
Seattle, WA 98195

Revised - November 8, 2000

## Abstract

We consider the Cauchy data associated to the Schrödinger equation with a potential on a bounded domain  $\Omega \subset \mathbb{R}^n, n \geq 3$ . We show that the integral of the potential over a two-plane  $\Pi$  is determined by the Cauchy data of certain exponentially growing solutions on any open subset  $\mathcal{U} \subset \partial\Omega$  which contains  $\Pi \cap \partial\Omega$ .

## 0 Introduction

For  $\Omega$  a bounded domain in  $\mathbb{R}^n$  with Lipschitz boundary,  $\partial\Omega$ , and real-valued  $q(x) \in L^\infty(\Omega)$ , let

$$(0.1) \quad \Lambda_q : H^{\frac{1}{2}}(\partial\Omega) \rightarrow H^{-\frac{1}{2}}(\partial\Omega)$$

be the Dirichlet-to-Neumann map associated with the operator  $\Delta + q$  on  $\Omega$ , which is defined if  $\lambda = 0$  is not a Dirichlet eigenvalue for  $\Delta + q$  on  $\Omega$ . More generally, one may consider the set of Cauchy data of solutions of

---

\*Partially supported by NSF grant DMS-9877101.

†Partially supported by NSF grant DMS-9705792 and the Royal Research Fund at the University of Washington

$(\Delta + q(x))v = 0$ , which is defined even if  $\lambda = 0$  is a Dirichlet eigenvalue. Set

(0.2)

$$\mathcal{CD}_q = \left\{ (v|_{\partial\Omega}, \frac{\partial v}{\partial n}|_{\partial\Omega}) \in H^{\frac{1}{2}}(\partial\Omega) \times H^{-\frac{1}{2}}(\partial\Omega) : v \in H^1(\Omega), (\Delta + q)v = 0 \right\},$$

which is a subspace of  $H^{\frac{1}{2}} \times H^{-\frac{1}{2}}$ ; if  $\Lambda_q$  is defined, then  $\mathcal{CD}_q$  is simply the graph of  $\Lambda_q$ .

This paper is concerned with the problem of obtaining partial knowledge of  $q(x)$  from partial knowledge of  $\mathcal{CD}_q$ , namely its restriction to certain “small” open subsets of the boundary. The approach taken here is to use concentrated, exponentially growing, approximate solutions to relate  $\mathcal{CD}_q$  on an open set  $\mathcal{U} \subset \partial\Omega$  to the two-plane transform of the potential  $q(x)$  on two-planes whose intersections with  $\partial\Omega$  are contained in  $\mathcal{U}$ .

Let  $M_{2,n}$  denote the  $(3n - 6)$ -dimensional Grassmannian of all affine two-planes  $\Pi \subset \mathbb{R}^n$ , and

$$(0.3) \quad R_{2,n}f(\Pi) = \int_{\Pi} f(y) d\lambda_{\Pi}(y), f \in L^2(\mathbb{R}^n),$$

denote the two-plane transform on  $\mathbb{R}^n$  [H65, H80]. Here,  $d\lambda_{\Pi}$  is two-dimensional Lebesgue measure on  $\Pi \in M_{2,n}$ , which can be defined by

$$(0.4) \quad \langle f, d\lambda_{\Pi} \rangle = \lim_{\epsilon \rightarrow 0} \frac{1}{|B^{n-2}(0; \epsilon)|} \int_{\{dist(x, \Pi) < \epsilon\}} f(x) dx.$$

(Note that for  $n = 3$ ,  $R_{2,3}$  is just the usual Radon transform on  $\mathbb{R}^3$ .) We will also need the variant of  $d\lambda_{\Pi}$  defined relative to  $\Omega$ :

$$(0.5) \quad \langle f, d\lambda_{\Pi}^{\Omega} \rangle = \lim_{\epsilon \rightarrow 0} \frac{1}{|B^{n-2}(0; \epsilon)|} \int_{\Omega \cap \{dist(x, \Pi) < \epsilon\}} f(x) dx,$$

which gives rise to a two-plane transform relative to  $\Omega$ ,

$$(0.6) \quad R_{2,n}^{\Omega}f(\Pi) = \int_{\Pi} f(x) d\lambda_{\Pi}^{\Omega}(x).$$

Note that if  $\partial\Omega$  is  $C^1$  and  $\Pi \cap \partial\Omega$  transversally, then  $\langle d\lambda_{\Pi}^{\Omega}, f \rangle = \langle d\lambda_{\Pi}, f \cdot \chi_{\Omega} \rangle$  and  $R_{2,n}^{\Omega}f(\Pi) = R_{2,n}(f \cdot \chi_{\Omega})(\Pi)$ .

For each choice of an orthonormal basis for  $\Pi_0$ , the translate of  $\Pi$  passing through the origin, as well as other arbitrary choices made below, we will construct a family,  $\mathcal{F}_q = \{v_z(x) : z \in \mathbb{C}, |z| \geq C\}$ , of exponentially growing

solutions of  $(\Delta + q(x))v = 0$ , concentrated near  $\Pi$ . Using these families, we formulate

**Definition** (i) If  $\mathcal{U} \subset \partial\Omega$  is open,  $\mathcal{CD}_{q_1}$  and  $\mathcal{CD}_{q_2}$  are *equal on  $\mathcal{U}$  relative to  $\mathcal{F}$*  at  $z \in \mathbb{C}$  if the solutions in  $\mathcal{F}_{q_1}$  and  $\mathcal{F}_{q_2}$  corresponding to opposite exponential growths,  $v_z^{(1)}$  and  $v_{-z}^{(2)}$ , have the same Cauchy data on  $\mathcal{U}$ :

$$(v_z^{(1)}|_{\mathcal{U}}, \frac{\partial v_z^{(1)}}{\partial n}|_{\mathcal{U}}) = (v_{-z}^{(2)}|_{\mathcal{U}}, \frac{\partial v_{-z}^{(2)}}{\partial n}|_{\mathcal{U}}).$$

(ii)  $\mathcal{CD}_{q_1}$  and  $\mathcal{CD}_{q_2}$  are *equal on  $\mathcal{U}$  for a sequence of exponentially growing solutions* if  $\mathcal{CD}_{q_1}$  and  $\mathcal{CD}_{q_2}$  are equal on  $\mathcal{U}$  relative to  $\mathcal{F}$  at  $z = z_j$  for some sequence  $\{z_j\}_1^\infty \subset \mathbb{C}$  with  $|z_j| \rightarrow \infty$ .

We may now state the main result proved here. For each  $\Pi \in M_{2,n}$ , let  $\gamma_\Pi = \Pi \cap \partial\Omega \subset \partial\Omega$ , and let  $H^s(\Omega)$  denote the standard Sobolev space of distributions with  $s$  derivatives in  $L^2(\Omega)$ .

**Theorem 1** *Let  $n \geq 3$ . Assume  $\partial\Omega$  is Lipschitz and potentials  $q_1(x)$  and  $q_2(x)$  are in  $H^s(\Omega)$ , for some  $s > \frac{n}{2}$ . Let  $\Pi \in M_{2,n}$  and  $\mathcal{F}_{q_1}$  and  $\mathcal{F}_{q_2}$  be families of exponentially growing solutions associated to  $q_1$  and  $q_2$ . If, for some fixed neighborhood  $\mathcal{U}_\Pi$  of  $\gamma_\Pi$  in  $\partial\Omega$ ,  $\mathcal{CD}_{q_1}$  and  $\mathcal{CD}_{q_2}$  are equal on  $\mathcal{U}_\Pi$  for a sequence of exponentially growing solutions, then*

$$(0.7) \quad R_{2,n}^\Omega(q_1 - q_2)(\Pi) = 0,$$

$$i.e., \int q_1(y) d\lambda_\Pi^\Omega(y) = \int q_2(y) d\lambda_\Pi^\Omega(y).$$

If  $\mathcal{CD}_{q_1}$  and  $\mathcal{CD}_{q_2}$  equal on all of  $\partial\Omega$  relative to  $\mathcal{F}$ , then this implies that  $R_{2,n}((q_1 - q_2)\chi_\Omega)(\Pi) = 0$ ,  $\forall \Pi \in M_{2,n}$ , which by the uniqueness theorem for  $R_{2,n}$  yields that  $q_1 - q_2 \equiv 0$  on  $\Omega$ , providing a variant of the global uniqueness theorem for the Dirichlet-to-Neumann map [SU87a]. (We note that our technique is limited to three or more dimensions and says nothing in the case  $n = 2$  [N96].) However, one is also able to obtain *local* uniqueness results by replacing the uniqueness theorem for the two-plane transform with Helgason's support theorem [H80, Cor. 2.8]: if  $C \subset \mathbb{R}^n$  is a closed, convex set and  $f(x)$  a function<sup>1</sup> such that  $R_{2,n}f(\Pi) = 0$  for all  $\Pi$  disjoint from  $C$ , then  $\text{supp}(f) \subset C$ . We then immediately obtain the following two results.

---

<sup>1</sup>The support and uniqueness theorems are usually stated under the assumption that  $f(x)$  is continuous, of rapid decay in the case of the support theorem, but the proofs in [H80] are easily seen to extend to the case where  $f(x) = q(x)\chi_\Omega(x)$  with  $\Omega \subset \mathbb{R}^n$  bounded,  $q \in C(\overline{\Omega})$ .

**Theorem 2** Suppose  $\partial\Omega$  and potentials  $q_1, q_2$  are as in Thm. 1., and  $C \subset \Omega$  is a closed, convex set. If, for all  $\Pi \in M_{2,n}$  such that  $\Pi \cap C = \emptyset$ , there is some neighborhood  $\mathcal{U}_\Pi$  of  $\gamma_\Pi$  on which  $\mathcal{CD}_{q_1}$  and  $\mathcal{CD}_{q_2}$  are equal for some sequence of exponentially growing solutions, then  $\text{supp}(q_1 - q_2) \subseteq C$ , i.e.,  $q_1 = q_2$  on  $\Omega \setminus C$ .

**Theorem 3** Suppose  $\partial\Omega$  is  $C^2$  and strictly convex, and potentials  $q_1, q_2$  are as in Thm. 1. If, for some  $r > 0$ ,  $\mathcal{CD}_{q_1}$  and  $\mathcal{CD}_{q_2}$  are equal on  $B$  for some sequence of exponentially growing solutions for all surface balls  $B = B^n(x_0; r) \cap \partial\Omega \subset \partial\Omega$ , then

$$\text{dist}(\text{supp}(q_1 - q_2), \partial\Omega) \geq Cr^2,$$

i.e.,  $q_1 = q_2$  on the tubular neighborhood  $\{x \in \overline{\Omega} : \text{dist}(x, \partial\Omega) \leq Cr^2\}$  of  $\partial\Omega$  in  $\overline{\Omega}$ .

**Remark**

The conclusions of Thms. 2 and 3 can be strengthened by combining them with a result in Isakov [Is]. Namely, if either  $C \subset\subset \Omega$  in Thm. 2, or the assumption of Thm. 3 holds for some  $r > 0$ , we can conclude from Thm. 2 or 3 that  $\text{supp}(q_1 - q_2) \subset\subset \Omega$ . By Ex. 5.7.4 in [Is], based on a technique of Kohn and Vogelius [KV85], this, together with the condition that  $\Lambda_{q_1} = \Lambda_{q_2}$  on some open set  $\mathcal{U} \subset \partial\Omega$ , implies that  $q_1 \equiv q_2$  everywhere on  $\Omega$ . We are indebted to Adrian Nachman for pointing this out to us.

The authors would like to thank Alexander Bukhgeim and Masaru Ikehata for pointing out errors in an earlier version of this paper.

## 1 Approximate solutions

To prove Thm. 1, we first construct exponentially growing *approximate* solutions for  $(\Delta + q)v = 0$ . As considered in [C, SU86, SU87a], let

$$\mathcal{Q} = \{\rho \in \mathbb{C}^n : \rho \cdot \rho = 0\}$$

be the (complex) characteristic variety of  $\Delta$ . Each  $\rho \in \mathcal{Q}$  can be written as  $\rho = |\rho| \frac{\rho}{|\rho|} = \frac{1}{\sqrt{2}} |\rho| (\omega_R + i\omega_I) \in \mathbb{R} \cdot (S^{n-1} + iS^{n-1})$ , with  $\omega_R \cdot \omega_I = 0$ . For  $\rho \in \mathcal{Q}$ , let  $\Delta_\rho = \Delta + 2\rho \cdot \nabla$ . Then

$$(1.1) \quad \Delta_\rho + q(x) = e^{-\rho \cdot x} (\Delta + q(x)) e^{\rho \cdot x},$$

so that, with  $v(x) = e^{\rho \cdot x} u(x)$ ,

$$(1.2) \quad (\Delta_\rho + q(x))u(x) = w(x) \Leftrightarrow (\Delta + q(x))v(x) = e^{\rho \cdot x} w(x)$$

and, in particular,  $(\Delta_\rho + q(x))u(x) = 0 \Leftrightarrow (\Delta + q(x))v(x) = 0$ .

Now, given a potential  $q(x)$  and a two-plane  $\Pi \in M_{2,n}$ , we will construct an approximate solution  $u_{app}$  to  $(\Delta_\rho + q)u = 0$ , supported near  $\Pi$ :

**Theorem 4** *Let  $\Omega$  be Lipschitz and  $q(x) \in H^s(\Omega)$  for some  $s > \frac{n}{2}$ . Then, for any  $0 < \beta < \frac{1}{4}$  fixed, the following holds:  $\exists \epsilon > 0$  such that, for any  $\rho = \frac{1}{\sqrt{2}}|\rho|(\omega_R + i\omega_I) \in \mathcal{Q}$  and any two-plane  $\Pi$  parallel to  $\Pi_0 = \text{span}\{\omega_R, \omega_I\}$ , we can find an approximate solution  $u_{app} = u_{app}(x, \rho, \Pi)$  to  $(\Delta_\rho + q(x))u = 0$  satisfying*

$$(1.5) \quad \|u_{app}\|_{L^2(\mathbb{R}^n)} \leq C, \quad \|u_{app}\|_{L^2(\Omega)} \simeq [\lambda_\Pi^\Omega(\Pi \cap \Omega)]^{\frac{1}{2}} \text{ as } |\rho| \rightarrow \infty$$

$$(1.6) \quad \text{supp}(u_{app}) \subset \left\{ x \in \mathbb{R}^n : \text{dist}(x, \Pi) \leq \frac{2}{|\rho|^\beta} \right\}$$

and

$$(1.7) \quad \|(\Delta_\rho + q)u_{app}\|_{L^2(\mathbb{R}^n)} \leq \frac{C_\epsilon}{|\rho|^\epsilon}.$$

Furthermore, for any two such solutions,  $u_{app}^{(1)}, u_{app}^{(2)}$ , associated with possibly different potentials  $q_1(x), q_2(x)$  and with  $\rho_1 \in \mathcal{Q}, \rho_2 = e^{i\theta}\rho_1$  or  $\rho_2 = e^{i\theta}\overline{\rho_1} \in \mathcal{Q}$ ,

$$(1.8) \quad u_{app}^{(1)}(\cdot, \rho_1, \Pi)u_{app}^{(2)}(\cdot, \rho_2, \Pi) \rightarrow d\lambda_\Pi^\Omega \text{ weakly as } |\rho_1| \rightarrow \infty.$$

In fact, as will be seen below,  $u_{app} = u_0 + u_1$  with  $u_0$  depending only on  $\Pi$  and  $|\rho|$  and satisfying (1.5).

Now, we may apply the results of [SU86, SU87a] (see also [Ha96]) to find a solution  $u_2$  of

$$(\Delta_\rho + q)u_2 = -(\Delta_\rho + q)u_{app} \in L_{comp}^2(\mathbb{R}^n),$$

uniformly in  $H_t^1$  and with a gain of  $|\rho|^{-1}$  in  $L_t^2$ , as long as  $|\rho| \geq C$  with  $C$  depending only on  $\|q\|_\infty$  and  $\text{diam}(\Omega)$ . Here,  $H_t^s$  and  $L_t^2$  the weighted versions of these spaces, as in [SU87a], for some fixed  $-1 < t < 0$ . By these results and (1.7),

$$\|u_2\|_{H_t^1(\mathbb{R}^n)} \leq c\|(\Delta_\rho + q)u_{app}\|_{L_{t+1}^2(\mathbb{R}^n)} \leq c|\rho|^{-\epsilon}, \quad \|u_2\|_{L_t^2} \leq C|\rho|^{-1-\epsilon}.$$

(The statements in [SU86,SU87a] are for  $q \in C^\infty$ , but the proofs are easily seen to hold if  $q \in H^s(\Omega)$  with  $s > \frac{n}{2}$ . Also, the weights will be irrelevant since we will be working on  $\Omega$ .) Thus,  $u = u_{app} + u_2 = u_0 + u_1 + u_2$  is an exact solution of  $(\Delta_\rho + q)u = 0$  on  $\mathbb{R}^n$ , satisfying

$$\|u - u_0\|_{L^2} \leq c|\rho|^{-\epsilon} \text{ and } \|u_2\|_{H^s} \leq |\rho|^{s-1-\epsilon}, \forall 0 \leq s \leq 1.$$

Finally,

$$\mathcal{F}_q = \{v_z : |z| \geq C\} = \{e^{\rho \cdot x} u(x, \Pi, \rho) : \rho = \operatorname{Re}(z)\omega_R + i\operatorname{Im}(z)\omega_I, |z| \geq C\}$$

is the associated family of exponentially growing solutions used in the statements of the theorems. To prove Thm. 1, we assume that  $q_1, q_2$  and  $\Pi \in M_{2,n}$ ,

$\mathcal{U}_\Pi \subset \partial\Omega$  are as in its statement. We will make use of a variant of Alessandrini's identity [A]. For  $j = 1, 2$ , let  $v_{\rho_j}^{(j)}$  be the exact solution to  $(\Delta + q_j)v = 0$  constructed above, so that  $v_{\rho_j}^{(j)}(x) = e^{\rho_j \cdot x} u^{(j)}(x, \Pi, \rho_j)$ , with  $u^{(j)} = u_{app}^{(j)} + u_2^{(j)}$ . Taking  $\rho_1 = \rho, \rho_2 = -\rho$ , consider the quantity

$$I = \int_{\partial\Omega} \frac{\partial v_\rho^{(1)}}{\partial n} \cdot v_{-\rho}^{(2)} - v_\rho^{(1)} \cdot \frac{\partial v_{-\rho}^{(2)}}{\partial n} d\sigma.$$

Under the assumption that  $v_\rho^{(1)}$  and  $v_{-\rho}^{(2)}$  have the same Cauchy data on  $\mathcal{U}_\Pi$ ,  $I$  is equal to the integral of the same expression over  $\partial\Omega \setminus \mathcal{U}_\Pi$ . Observing that

$$\frac{\partial v_\rho^{(1)}}{\partial n} = e^{\rho \cdot x} \left( \frac{\partial}{\partial n} + (\rho \cdot n(x)) \right) u^{(1)} \text{ and } \frac{\partial v_{-\rho}^{(2)}}{\partial n} = e^{-\rho \cdot x} \left( \frac{\partial}{\partial n} - (\rho \cdot n(x)) \right) u^{(2)},$$

we see that the exponentials cancel and the integrand of  $I$  is

$$= \frac{\partial u^{(1)}}{\partial n} \cdot u^{(2)} - u^{(1)} \cdot \frac{\partial u^{(2)}}{\partial n} + 2(\rho \cdot n(x)) u^{(1)} u^{(2)}.$$

Since (1.6) implies that  $\operatorname{supp}(u_{app}^{(j)}|_{\partial\Omega}), \operatorname{supp}(\frac{\partial u_{app}^{(j)}}{\partial n}|_{\partial\Omega}) \subset \mathcal{U}_\Pi$  for  $|\rho|$  sufficiently large, we have that

$$I = \int_{\partial\Omega \setminus \mathcal{U}_\Pi} \frac{\partial u_2^{(1)}}{\partial n} \cdot u_2^{(2)} - u_2^{(1)} \cdot \frac{\partial u_2^{(2)}}{\partial n} + 2(\rho \cdot n(x)) u_2^{(1)} u_2^{(2)} d\sigma.$$

We estimate

$$\begin{aligned}
\left| \int_{\partial\Omega \setminus \mathcal{U}_\Pi} \frac{\partial u_2^{(1)}}{\partial n} \cdot u_2^{(2)} d\sigma \right| &\leq \left\| \frac{\partial u_2^{(1)}}{\partial n} \right\|_{H^{-\frac{1}{2}}(\partial\Omega)} \cdot \|u_2^{(2)}\|_{H^{\frac{1}{2}}(\partial\Omega)} \\
&\leq \|u_2^{(1)}\|_{H^{\frac{1}{2}}(\partial\Omega)} \cdot \|u_2^{(2)}\|_{H^{\frac{1}{2}}(\partial\Omega)} \\
&\leq C \|u_2^{(1)}\|_{H^1(\Omega)} \cdot \|u_2^{(2)}\|_{H^1(\Omega)} \text{ by Sobolev restriction} \\
&\leq C \|u_2^{(1)}\|_{H_t^1(\mathbb{R}^n)} \cdot \|u_2^{(2)}\|_{H_t^1(\mathbb{R}^n)} \text{ since } \Omega \text{ compact} \\
&\leq C |\rho|^{-2\epsilon} \rightarrow 0 \text{ as } |\rho| \rightarrow \infty
\end{aligned}$$

and similarly for the second term. Now note that  $|\rho \cdot n(x)| \leq c|\rho|$  since  $\partial\Omega$  is Lipschitz, and

$$\|u_2^{(j)}\|_{L^2(\partial\Omega)} \leq \|u_2^{(j)}\|_{H^\sigma(\partial\Omega)} \leq c_\sigma \|u_2^{(j)}\|_{H^{\sigma+\frac{1}{2}}(\Omega)} \leq c'_\sigma |\rho|^{\sigma-\frac{1}{2}-\epsilon}$$

for any  $\sigma > 0$ , and thus the third term is dominated by  $(c'_\sigma)^2 |\rho| \cdot |\rho|^{2\sigma-1-2\epsilon} \rightarrow 0$  as  $|\rho| \rightarrow 0$  if we choose  $0 < \sigma < \epsilon$ .

On the other hand,

$$\begin{aligned}
I &= \int_{\partial\Omega} \frac{\partial v^{(1)}}{\partial n} \cdot v^{(2)} - v^{(1)} \cdot \frac{\partial v^{(2)}}{\partial n} d\sigma \\
&= \int_{\Omega} \Delta(v^{(1)}) \cdot v^{(2)} - v^{(1)} \cdot \Delta(v^{(2)}) dx \text{ by Green's Thm.} \\
&= \int_{\Omega} (-q_1 v^{(1)}) \cdot v^{(2)} - v^{(1)} \cdot (-q_2 v^{(2)}) dx \\
&= \int_{\Omega} (q_2 - q_1) v^{(1)} v^{(2)} dx = \int_{\Omega} (q_2 - q_1) u^{(1)} u^{(2)} dx
\end{aligned}$$

since the exponentials cancel. As  $u^{(1)} \cdot u^{(2)} = (u_{app}^{(1)} + u_2^{(1)}) \cdot (u_{app}^{(2)} + u_2^{(2)})$  and the leading term  $u_{app}^{(1)} u_{app}^{(2)} \rightarrow d\lambda_\Pi^\Omega$  weakly as  $|\rho| \rightarrow \infty$  by (1.8), while the remaining terms  $\rightarrow 0$  since  $\|u_{app}^{(j)}\|_{L^2(\Omega)} \leq C$  by (1.5) and  $\|u_2^{(j)}\|_{L^2(\Omega)} \leq c|\rho|^{-1-\epsilon}$ , we conclude that  $I \rightarrow R_{2,n}^\Omega(q_2 - q_1)(\Pi)$  as  $|\rho| \rightarrow \infty$ , finishing the proof of Thm. 1.

Now, to start the proof of Thm. 4 we may use the rotation invariance of  $\Delta$  and the invariance of  $\mathcal{Q}$  under  $S^1 = \{e^{i\theta}\}$ , and note that it suffices to treat the case<sup>2</sup>  $\rho = |\rho|(\vec{e}_1 + i\vec{e}_2)$ , where  $\{\vec{e}_1, \dots, \vec{e}_n\}$  is the standard orthonormal

---

<sup>2</sup>Of course, the length of this element of  $\mathcal{Q}$  is  $\sqrt{2}|\rho|$ , but this is irrelevant for the proofs, and denoting the length of  $|\rho|(\vec{e}_1 + i\vec{e}_2)$  by  $|\rho|$  is notationally convenient.

basis for  $\mathbb{R}^n$ . Write  $x \in \mathbb{R}^n$  as  $x = (x', x'') \in \mathbb{R}^2 \times \mathbb{R}^{n-2}$  and similarly  $\xi = (\xi', \xi'')$ .

If  $\Pi \in M_{2,n}$  is parallel to  $\text{span}\{\omega_R, \omega_I\} = \text{span}\{\vec{e}_1, \vec{e}_2\} = \mathbb{R}^2 \times \{0\}$ , then  $\Pi = \text{span}\{\vec{e}_1, \vec{e}_2\} + (0, x''_0)$  for some  $x''_0 \in \mathbb{R}^{n-2}$ . Given  $|\rho| > 1$  and  $x''_0 \in \mathbb{R}^{n-2}$ , we will define an approximate solution  $u(x, \rho, \Pi)$  to  $(\Delta_\rho + q(x))u = 0$  on  $\mathbb{R}^n$ , of the form  $u(x, \rho, \Pi) = u_0(x, \rho, \Pi) + u_1(x, \rho, \Pi)$ .

For notational convenience, we will usually suppress the dependence on  $\rho$  and  $\Pi$  and simply write  $u(x) = u_0(x) + u_1(x)$ . We will use various cutoff functions  $\chi_j$ ; for  $j$  even or odd,  $\chi_j$  will always denote a function of  $x'$  or  $x''$ , respectively. Also,  $B^m(a; r)$  and  $S^{m-1}(a; r)$  will denote the closed ball and sphere of radius  $r$  centered at a point  $a \in \mathbb{R}^m$ .

To define  $u_0$ , first fix  $\chi_0 \in C_0^\infty(\mathbb{R}^2)$  with  $\chi_0 \equiv 1$  on  $B^2(0; R)$  for any  $R > \sup\{|x'| : (x', x'') \in \Omega \text{ for some } x'' \in \mathbb{R}^{n-2}\}$ ; let  $C_0 = \|\chi_0\|_{L^2(\mathbb{R}^2)}$ . Secondly, let  $\psi_1 \in C_0^\infty(\mathbb{R}^{n-2})$  be radial, nonnegative, supported in the unit ball, and satisfy

$$\int_{\mathbb{R}^{n-2}} (\psi_1(x''))^2 dx'' = 1.$$

Now, for  $\beta > 0$  to be fixed later, we let  $\delta$  be the small parameter  $\delta = |\rho|^{-\beta}$  and define

$$\chi_1(x'') = \delta^{-\frac{n-2}{2}} \psi_1\left(\frac{x' - x''_0}{\delta}\right),$$

so that

$$(1.9) \quad \|\chi_1\|_{L^2(\mathbb{R}^{n-2})} = \|\psi_1\|_{L^2(\mathbb{R}^{n-2})} = 1, \quad \forall \delta > 0.$$

Set  $u_0(x) = u_0(x', x'') = \chi_0(x')\chi_1(x'')$ ; then  $u_0$  is real,  $\|u_0\|_{L^2(\mathbb{R}^n)} = C_0$  and  $\|u_0\|_{L^2(\Omega)} \rightarrow [\lambda_\Pi(\Pi \cap \Omega)]^{\frac{1}{2}}$  as  $\delta \rightarrow 0^+$ , i.e., as  $|\rho| \rightarrow \infty$ . Note also that  $\|u_0\|_{H^1} \leq c\delta^{-1} = c|\rho|^\beta$ , so that  $\|u_0\|_{H^s} \leq c|\rho|^{s\beta}$  for  $0 \leq s \leq 1$ . Since  $\Delta_\rho = \Delta + 2\rho \cdot \nabla = \Delta + 2|\rho|(\vec{e}_1 + i\vec{e}_2) \cdot \nabla = \Delta + 4|\rho|\bar{\partial}_{x'}$  and  $\rho \perp \mathbb{R}^{n-2}$ ,

$$\begin{aligned} (\Delta_\rho + q(x))u_0 &= (\Delta\chi_0) \cdot \chi_1 + 2(\nabla\chi_0) \cdot (\nabla\chi_1) + \chi_0(\Delta\chi_1) \\ &\quad + 2(\rho \cdot \nabla)(\chi_0)\chi_1 + 2\chi_0(\rho \cdot \nabla)(\chi_1) + q\chi_0\chi_1 \\ &= \chi_0(x')(\Delta_{x''} + q)(\chi_1)(x'') \text{ on } B^2(0; R) \times \mathbb{R}^{n-2}, \end{aligned}$$

the first and fourth terms after the first equality vanishing because  $(\rho \cdot \nabla)(\chi_0) = 2\bar{\partial}\chi_0 \equiv 0$  on  $B^2(0; R)$ , and the second and fifth equalling zero because  $\nabla\chi_1 \perp \mathbb{R}^2$ .



To define the second term in the approximate solution,  $u_1(x)$ , we make use of a truncated form of the Faddeev Green function,  $G_\rho$ , and an associated projection operator. The operator  $\Delta_\rho$  has, for  $\rho \in \mathcal{Q}$ , (full) symbol

$$(1.10) \quad \sigma(\xi) = -[(|\xi|^2 - 2|\rho|\omega_I \cdot \xi) + i2|\rho|(\omega_R \cdot \xi)],$$

and so for  $\frac{\rho}{|\rho|} = e_1 + ie_2$ , we have

$$\sigma(\xi) = -[(|\xi - |\rho|e_2|^2 - |\rho|^2) + i(2|\rho|\xi_1)],$$

which has (full) characteristic variety

$$(1.11) \quad \begin{aligned} \Sigma_\rho &= \{\xi \in \mathbb{R}^n : \xi_1 = 0, |\xi - |\rho|e_2| = |\rho|\} \\ &= \{0\} \times S^{n-2}(|\rho|, 0, \dots, 0; |\rho|) \subset \mathbb{R}_{\xi_1} \times \mathbb{R}_{\xi_2, \xi''}^{n-1}. \end{aligned}$$

The Faddeev Green function is then defined by  $G_\rho = (-\sigma(\xi)^{-1})^\vee \in \mathcal{S}'(\mathbb{R}^n)$ . We now introduce, for an  $\epsilon_0 > 0$  to be fixed later, a tubular neighborhood of  $\Sigma_\rho$ ,

$$(1.12) \quad T_\rho = \{\xi : \text{dist}(\xi, \Sigma_\rho) < |\rho|^{-\frac{1}{2}-\epsilon_0}\},$$

as well as its complement,  $T_\rho^C$ , and let  $\chi_{T_\rho}$ ,  $\chi_{T_\rho^C}$  be their characteristic functions. Define a projection operator,  $P_\rho$ , and a truncated Green function,  $\tilde{G}_\rho$ , by

$$(1.13) \quad \widehat{P_\rho f}(\xi) = \chi_{T_\rho}(\xi) \cdot \widehat{f}(\xi) \text{ and}$$

$$(1.14) \quad (\tilde{G}_\rho f)^\wedge(\xi) = \chi_{T_\rho^C}(\xi) \cdot [-\sigma(\xi)]^{-1} \widehat{f}(\xi)$$

for  $f \in \mathcal{S}(\mathbb{R}^n)$ . Note that  $\Delta_\rho \tilde{G}_\rho = I - P_\rho$ .

Choose a  $\psi_3 \in C_0^\infty(\mathbb{R}^{n-2})$ , supported in  $B^{n-2}(0; 2)$ , radial and with  $\psi_3 \equiv 1$  on  $\text{supp}(\psi_1)$ , and set  $\chi_3(x'') = \psi_3(\frac{x'' - x_0''}{\delta})$ . We now define the second term,  $u_1(x, \rho, \Pi)$  in the approximate solution by

$$(1.15) \quad u_1(x) = -\chi_3(x'') \tilde{G}_\rho((\Delta_\rho + q(x))u_0(x))$$

and set  $u(x) = u_0(x) + u_1(x)$ . Then  $u_1$  (as well as  $u_0$ ) is supported in  $\{x : \text{dist}(x, \Pi) \leq 2\delta\}$ , yielding (1.6). We will see below that  $\|u_1\|_{L^2(\Omega)} \leq C|\rho|^{-\epsilon}$  as  $|\rho| \rightarrow \infty$ , so that (1.5) holds as well, so that the first part of (1.9) holds

as well. To start the proof of (1.7), note that

$$\begin{aligned}
(\Delta_\rho + q)(u_0 + u_1) &= (\Delta_\rho + q)u_0 - (\Delta_\rho + q)\chi_3\tilde{G}_\rho((\Delta_\rho + q)u_0) \\
&= (\Delta_\rho + q)u_0 - \chi_3(\Delta_\rho + q)\tilde{G}_\rho((\Delta_\rho + q)u_0) \\
&\quad - [\Delta_\rho + q, \chi_3]\tilde{G}_\rho((\Delta_\rho + q)u_0) \\
&= (\Delta_\rho + q)u_0 - \chi_3(I - P_\rho)(\Delta_\rho + q)u_0 - \chi_3q\tilde{G}_\rho(\Delta_\rho + q)u_0 \\
&\quad - 2(\nabla\chi_3 \cdot \nabla_{x''})\tilde{G}_\rho(\Delta_\rho + q)u_0 - (\Delta_{x''}\chi_3)\tilde{G}_\rho(\Delta_\rho + q)u_0 \\
&= \chi_3P_\rho(\Delta_\rho + q)u_0 \\
&\quad - [q\chi_3 + 2(\nabla\chi_3 \cdot \nabla_{x''}) - (\Delta_{x''}\chi_3)]\tilde{G}_\rho(\Delta_\rho + q)u_0
\end{aligned}$$

on  $\Omega$ , since  $\chi_3 \equiv 1$  on  $\text{supp}(\chi_1)$ . Now, since  $q_1\chi_3 \in L^\infty$ ,  $|\nabla\chi_3| \leq C\delta^{-1} = c|\rho|^\beta$  and  $|\Delta_{x''}\chi_3| \leq C\delta^{-2} = c|\rho|^{2\beta}$ , (1.7) will follow if we can show that for some  $\epsilon > 0$ ,

$$(1.16) \quad \|P_\rho(\Delta_\rho + q)u_0\|_{L^2(\Omega)} \leq C|\rho|^{-\epsilon},$$

$$(1.17) \quad \| |D''|\tilde{G}_\rho(\Delta_\rho + q)u_0 \|_{L^2(\Omega)} \leq C|\rho|^{-\beta-\epsilon}, \text{ and}$$

$$(1.18) \quad \|\tilde{G}_\rho(\Delta_\rho + q)u_0\|_{L^2(\Omega)} \leq C|\rho|^{-2\beta-\epsilon},$$

with  $C$  independent of  $|\rho| > 1$ . Before proceeding to prove these, we note that for any  $u^{(1)}, u^{(2)}$  constructed in this way for the same two-plane  $\Pi$ ,

$$u_0^{(1)}(x)u_0^{(2)}(x) = \chi_0^2(x')\delta^{-(n-2)}\psi_1^2\left(\frac{x'' - x_0''}{\delta}\right) \rightarrow d\lambda_\Pi^\Omega \text{ in } \Omega$$

as  $\delta \rightarrow 0$  by (1.11), while  $u_1^{(1)}u_0^{(2)} + u_0^{(1)}u_1^{(2)} + u_1^{(1)}u_1^{(2)} \rightarrow 0$  in  $L^2(\Omega)$ , yielding (1.8). Thus, we are reduced to establishing (1.17–1.19).

## 2 $L^2$ estimates

We will first prove (1.17)–(1.19) under the simplifying assumption that  $q_1, q_2 \in C^{n-1+\sigma}(\overline{\Omega})$  for some  $\sigma > 0$ , turning to the Sobolev space case in Section 3. Start by noting that the desired estimates (1.17)–(1.19) cannot be simply obtained from operator norms; for example,  $\|P_\rho\|_{L^2 \rightarrow L^2} = 1$  for all  $\rho$ . One needs to make use of the special structure of  $(\Delta_\rho + q)u_0$ ; we first deal with  $\Delta_\rho u_0$ , leaving  $q(x) \cdot u_0$  for the end. So, we will show that  $\|P_\rho\Delta_\rho u_0\|_{L^2} \leq C|\rho|^{-\epsilon}$ , etc. Since  $\nabla\chi_0 \cdot \nabla\chi_1 \equiv 0$ ,

$$(2.1) \quad \Delta_\rho u_0 = \chi_0\Delta_{x''}\chi_1 + (\Delta_{x'} + 4|\rho|\bar{\partial}_{x'})(\chi_0) \cdot \chi_1.$$

The second term is supported on  $\Omega^c$ , but  $P_\rho$  and  $\tilde{G}_\rho$  are nonlocal operators and we need to control the contribution from this term. However, because  $\Delta_{x'}(\chi_0)$  is a fixed,  $\delta$ -independent element of  $C_0^\infty(\mathbb{R}^2)$ , this can be handled in the same way as the  $q(x) \cdot u_0$  terms of (1.17–1.19), which will be dealt with later. The contribution from  $4|\rho|\bar{\partial}\chi_0 \cdot \chi_1$  will be handled at the end.

So, for the time being, we are interested in estimating  $\|P_\rho(\chi_0(x')\Delta_{x''}\chi_1(x''))\|_{L^2}$ , etc. Now,  $\Delta_{x''}\chi_1(x'') = \delta^{-2}\chi_5(x'')$ , where  $\chi_5(x'') = \delta^{-\frac{n-2}{2}}\psi_5\left(\frac{x''-x_0''}{\delta}\right)$  is associated with the radial function  $\psi_5 = \Delta_{x''}\psi_1$  as  $\chi_1$  is associated with  $\psi_1$ . Note for future use that  $\widehat{\psi}_5$  vanishes to second order at 0. Of course,  $\chi_0 \in C_0^\infty \Rightarrow \widehat{\chi}_0 \in \mathcal{S}(\mathbb{R}^n)$ , but looking ahead to estimating the terms involving  $q(x) \cdot u_0(x)$ , we will now prove the analogues of (1.17–1.19) where  $P_\rho$  and  $\tilde{G}_\rho$  act on  $\chi_2(x')\Delta\chi_1(x'')$ , under the weaker assumption that  $\chi_2$  is radial and satisfies the uniform decay estimate

$$(2.2)_\alpha \quad |\widehat{\chi}_2(\xi)| \leq C(1 + |\xi|)^{-\alpha}$$

for some  $\alpha > 0$ .

Now, by (1.14) and Plancherel,

$$\begin{aligned} \|P_\rho(\chi_2\Delta\chi_1)\|_{L^2(\Omega)} &\leq \|(P_\rho(\chi_2\Delta\chi_1))^\wedge\|_{L^2(\mathbb{R}^n)} \\ &= \|\delta^{-2}|\widehat{\chi}_2(\xi')|\delta^{\frac{n-2}{2}}|\widehat{\psi}_5(\delta\xi'')|\|_{L^2(T_\rho)}. \end{aligned}$$

The characteristic variety  $\Sigma_\rho$ , of which  $T_\rho$  is a tubular neighborhood, passes through the origin, and we may represent  $\Sigma_\rho$  near  $O$  as a graph over the  $\xi''$ -plane:  $\Sigma_\rho = \Sigma_\rho^s \cup \Sigma_\rho^n \cup \Sigma_\rho^e$ , with

$$(2.3) \quad \begin{aligned} \Sigma_\rho^s &= \left\{ \xi_1 = 0, \xi_2 = |\rho| - (|\rho|^2 - |\xi''|^2)^{\frac{1}{2}}, |\xi''| \leq \frac{|\rho|}{2} \right\} \\ &\simeq \left\{ \xi_1 = 0, \xi_2 = \frac{|\xi''|}{2|\rho|}, |\xi''| \leq \frac{|\rho|}{2} \right\} \end{aligned}$$

a neighborhood of the south pole  $O$ ,

$$(2.4) \quad \begin{aligned} \Sigma_\rho^n &= \left\{ \xi_1 = 0, \xi_2 = |\rho| + (|\rho|^2 - |\xi''|^2)^{\frac{1}{2}}, |\xi''| \leq \frac{|\rho|}{2} \right\} \\ &\simeq \left\{ \xi_1 = 0, \xi_2 = 2|\rho| - \frac{|\xi''|^2}{2|\rho|}, |\xi''| \leq \frac{|\rho|}{2} \right\} \end{aligned}$$

a neighborhood of the north pole  $(0, 2|\rho|, 0, \dots, 0)$ , and  $\Sigma_\rho^e$  a neighborhood of the equator  $\{\xi \in \Sigma_\rho : \xi_2 = |\rho|\}$ . We have a corresponding decomposition

$T_\rho = T_\rho^s \cup T_\rho^n \cup T_\rho^e$ , where, e.g.,

$$(2.5) \quad T_\rho^s \simeq \left\{ (\xi', \xi'') : \xi' \in B^2 \left( \left( 0, \frac{|\xi''|^2}{2|\rho|} \right); |\rho|^{-\frac{1}{2}-\epsilon_0} \right), |\xi''| \leq \frac{|\rho|}{2} \right\}.$$

Recalling that  $\chi_2$  and  $\psi_3$  are radial, so are  $\widehat{\chi}_2$  and  $\widehat{\chi}_3$ , and by abuse of notation we consider these as functions of one variable satisfying  $(2.2)_\alpha$  and rapidly decreasing, respectively. Thus, using polar coordinates in  $\xi''$ ,

$$(2.6) \quad \begin{aligned} \|\widehat{\chi_2 \Delta \chi_1}\|_{L^2(T_\rho^s)}^2 &\simeq \int_0^{\frac{|\rho|}{2}} \int_{B^2((0, \frac{r^2}{2|\rho|}); |\rho|^{-\frac{1}{2}-\epsilon_0})} |\widehat{\chi}_2(\xi')|^2 d\xi' \delta^{n-6} |\widehat{\psi}_5(\delta r)|^2 r^{n-3} dr \\ &\simeq \int_0^{\sqrt{2}|\rho|^{\frac{1}{4}}} \int_{B^2((0,0); |\rho|^{-\frac{1}{2}-\epsilon_0})} |\widehat{\chi}_2|^2 d\xi' \delta^{n-6} |\widehat{\psi}_5(\delta r)|^2 r^{n-2} \frac{dr}{r} \\ &\quad + \int_{\sqrt{2}|\rho|^{\frac{1}{4}}}^{\frac{|\rho|}{2}} \left| \widehat{\chi}_2 \left( \frac{r^2}{2|\rho|} \right) \right|^2 \cdot |B^2((0,0); |\rho|^{-\frac{1}{2}})| \delta^{n-6} |\widehat{\psi}_5(\delta r)|^2 r^{n-2} \frac{dr}{r}. \end{aligned}$$

Since we will be taking  $\delta = |\rho|^{-\beta}$  with  $\beta < \frac{1}{4}$ , if we choose  $0 < \epsilon_0 < 2(\frac{1}{4} - \beta)$ , then the quantity  $|\rho|^{\frac{1}{4}}\delta \rightarrow \infty$  as  $|\rho| \rightarrow \infty$  and so

$$(2.7) \quad \begin{aligned} \|\widehat{\chi_2 \Delta \chi_1}\|_{L^2(T_\rho^s)}^2 &\leq c \frac{\delta^{-4}}{|\rho|^{1+2\epsilon_0}} \left( \int_0^{\sqrt{2}|\rho|^{\frac{1}{4}}\delta} |\widehat{\psi}_5(r)|^2 r^{n-2} \frac{dr}{r} \right) \\ &\quad + \int_{\sqrt{2}|\rho|^{\frac{1}{4}}\delta}^{\frac{|\rho|}{2}} \left| \widehat{\chi}_2 \left( \frac{r^2}{2\delta^2|\rho|} \right) \right|^2 |\widehat{\psi}_5(r)|^2 r^{n-2} \frac{dr}{r} \\ &\leq c(\delta^4|\rho|)^{-1}, \end{aligned}$$

which is  $\leq c|\rho|^{-2\epsilon}$  with  $\epsilon = \frac{1}{2}(1 - 4\beta) > 0$ .

The other contributions to  $\|P_\rho \chi_2 \Delta \chi_1\|_{L^2}$ , coming from  $T_\rho^n$  and  $T_\rho^e$  are handled similarly and are even smaller, due to the decrease of  $\widehat{\chi}_2$  and  $\widehat{\psi}_5$ .

We next turn to estimating  $\| |D''| \widetilde{G}_\rho \Delta_\rho u_0 \|_{L^2}$ ; by the remark above, we may concentrate on the  $\chi_2 \Delta \chi_1$  term of  $\Delta_\rho u_0$ . Then

$$(2.8) \quad \| |D''| \widetilde{G}_\rho (\chi_2 \Delta \chi_1) \|_{L^2(\Omega)}^2 \leq \| |\xi''| (\sigma(\xi))^{-1} (\chi_2 \Delta \chi_1)^\wedge(\xi) \|_{L^2(T_\rho^C)}^2.$$

We may cover  $T_\rho^C$  by  $T_\rho^{C,s} \cup T_\rho^{C,n} \cup T_\rho^{C,e} \cup T_\rho^{C,\infty}$ , where

$$(2.9) \quad T_\rho^{C,s} = \left\{ \xi : \xi' \in B^2 \left( \left( 0, \frac{|\xi''|^2}{2|\rho|} \right); |\rho|^{-\frac{1}{2}-\epsilon_0} \right)^C \cap B^2 \left( \left( 0, 2|\rho| - \frac{|\xi''|^2}{2|\rho|} \right); \frac{1}{4}|\rho| \right)^C, |\xi''| \leq \frac{|\rho|}{2} \right\},$$

$T_\rho^{C,n}$  is defined similarly,

$$(2.10) \quad T_\rho^{C,e} = \left\{ \xi : \frac{|\rho|}{4} < \xi_2 < \frac{7|\rho|}{4}, |\rho|^{-\frac{1}{2}} < \text{dist}(\xi, \Sigma_\rho) < |\rho|, |\xi''| < 2|\rho| \right\}$$

and

$$(2.11) \quad T_\rho^{C,\infty} = \left\{ \xi : |\xi| \geq 3|\rho|, |\xi''| \geq \frac{3}{2}|\rho| \right\}.$$

One has the lower bounds on  $\sigma$ ,

$$(2.12) \quad |\sigma(\xi)| \geq \begin{cases} C|\rho|\text{dist}(\xi, \Sigma_\rho), & |\xi| \leq 3|\rho| \\ C|\xi|^2, & |\xi| \geq 3|\rho| \end{cases}$$

with  $C$  (as always) uniform in  $|\rho|$ . The first inequality in (2.12) follows from noting that  $\frac{1}{2}\nabla\sigma(\xi) = (\xi - |\rho|\vec{e}_2) + i(|\rho|\vec{e}_1)$ , so that  $|\nabla\sigma(\xi)| = 2\sqrt{2}|\rho|$  on  $\Sigma_\rho$ , while the second follows from  $\text{Re}(\sigma(\xi)) = \text{dist}(\xi, |\rho|\vec{e}_2)^2 - |\rho|^2$ . Using the first estimate in (2.12), we can then dominate the contribution to the right side of (2.8) from the region  $T_\rho^{C,s}$  by

$$(2.13) \quad \delta^{n-6} \int_{|\xi''| \leq \frac{|\rho|}{2}} \int_{B^2\left((0, \frac{|\xi''|^2}{2|\rho|}); |\rho|^{-\frac{1}{2}-\epsilon_0}\right)^C} |\rho|^{-2} \left| \xi' - \frac{|\xi''|^2}{2|\rho|} \vec{e}_2 \right|^{-2} |\widehat{\chi}_2(\xi')|^2 d\xi' |\xi''|^2 |\widehat{\psi}_5(\delta\xi'')|^2 d\xi''.$$

The inner integral is the convolution

$$|\rho|^{-2} \left( |\widehat{\chi}_2|^2 *_{\mathbb{R}^2} \frac{\chi\{|\xi'| \geq |\rho|^{-\frac{1}{2}-\epsilon_0}\}}{|\xi'|^2} \right) \Big|_{\xi' = \frac{|\xi''|^2}{2|\rho|} \vec{e}_2}.$$

An elementary calculation shows that, for  $\widehat{\chi}_2$  satisfying (2.2) $_\alpha$  for some  $0 < \alpha < 1$ , and any  $0 < a < 1$ ,

$$(2.14) \quad |\widehat{\chi}_2|^2 *_{\mathbb{R}^2} \frac{\chi\{|\xi'| \geq a\}}{|\xi'|^2} \leq \begin{cases} C_1(1 + \log(a^{-1})), & |\xi'| \leq 1 \\ C_2|\xi'|^{-2} + C_3|\xi'|^{-2\alpha} \log\left(\frac{|\xi'|}{a}\right), & |\xi'| \geq 1, \end{cases}$$

so that, taking  $a = |\rho|^{-\frac{1}{2}-\epsilon_0}$  and  $|\xi'| = \frac{|\xi''|^2}{2|\rho|}$ , the inner integral in (2.13) is

$$\leq \begin{cases} C_1|\rho|^{-2} \log|\rho|, & 0 < |\xi''| \leq \sqrt{2}|\rho|^{\frac{1}{2}} \\ C_2|\xi''|^{-4} + C_3|\rho|^{2\alpha-2} |\xi''|^{-4\alpha} \log\left(\frac{|\xi''|^2}{2|\rho|^{\frac{1}{2}-\epsilon_0}}\right), & \sqrt{2}|\rho|^{\frac{1}{2}} \leq |\xi''| \leq \frac{|\rho|}{2}. \end{cases}$$

Employing polar coordinates in  $\xi''$  and rescaling by  $\delta$ , we see that (2.13) is

$$\begin{aligned} &\leq C_1 \delta^{-6} |\rho|^{-2} \log |\rho| \int_0^{\sqrt{2}|\rho|^{\frac{1}{2}}\delta} |\widehat{\psi}_5(r)|^2 r^n \frac{dr}{r} \\ &\quad + C_2 \delta^{-2} \int_{\sqrt{2}|\rho|^{\frac{1}{2}}\delta}^{\frac{|\rho|}{2}\delta} |\widehat{\psi}_5(r)|^2 r^{n-4} \frac{dr}{r} \\ &\quad + C_3 \delta^{4\alpha-4} |\rho|^{2\alpha-2} \log |\rho| \int_{\sqrt{2}|\rho|^{\frac{1}{2}}\delta}^{\frac{|\rho|}{2}\delta} |\widehat{\psi}_5(r)|^2 r^{n-2-4\alpha} \frac{dr}{r}. \end{aligned}$$

With  $\delta = |\rho|^{-\beta}$ ,  $\beta < \frac{1}{4}$ ,  $|\rho|^{\frac{1}{2}}\delta \rightarrow \infty$  as  $|\rho| \rightarrow \infty$ , and thus we estimate this for any  $N > 0$  (using the rapid decay of  $\widehat{\psi}_5$ ) by

$$C_1 |\rho|^{6\beta-2} \log |\rho| + C_2 \delta^{-2} (|\rho|^{\frac{1}{2}}\delta)^{-N} + C_3 |\rho|^{(4-4\alpha)\beta+2\alpha-2} \log |\rho| (|\rho|^{\frac{1}{2}}\delta)^{-N},$$

the first term of which will be less than the desired  $|\rho|^{-2\beta-2\epsilon}$ , for any  $\alpha > 0$ , if  $\beta < \frac{1}{4}$  and  $\epsilon = \frac{1}{2}(1-4\beta)$ ; the second and third terms are rapidly decaying simply because  $\beta < \frac{1}{2}$ .

Moving ahead for the moment to (1.18), the contribution to  $\|\widetilde{G}_\rho \chi_2 \Delta \chi_1\|_{L^2}^2$  (which we want  $\leq C|\rho|^{-4\beta-2\epsilon}$ ) from  $T_\rho^{C,s}$  is handled in the same fashion, the only differences being the absence of the multiplier  $|D''|^\wedge = |\xi''|$  on the left and the improved gain we are demanding on the right. Taking these into account, we need to control

$$\begin{aligned} (2.15) \quad &C_1 \delta^{-4} |\rho|^{-2} \log |\rho| \int_0^{\sqrt{2}|\rho|^{\frac{1}{2}}\delta} |\widehat{\psi}_5(r)|^2 r^{n-2} \frac{dr}{r} \\ &\quad + C_2 \int_{\sqrt{2}|\rho|^{\frac{1}{2}}\delta}^{\frac{1}{2}|\rho|\delta} |\widehat{\psi}_5(r)|^2 r^{n-6} \frac{dr}{r} \\ &\quad + C_3 \delta^{4\alpha-2} |\rho|^{2\alpha-2} \log |\rho| \int_{\sqrt{2}|\rho|^{\frac{1}{2}}\delta}^{\frac{1}{2}|\rho|\delta} |\widehat{\psi}_5(r)|^2 r^{n-4-4\alpha} \frac{dr}{r} \\ &\leq C_1 \delta^{-4} |\rho|^{-2} \log |\rho| + C_2 (|\rho|^{\frac{1}{2}}\delta)^{-N} + C_N \delta^{4\alpha-2} |\rho|^{2\alpha-2} \log |\rho| (|\rho|^{\frac{1}{2}}\delta)^{-N}, \end{aligned}$$

and this is  $\leq C|\rho|^{-4\beta-2\epsilon}$  provided  $\beta < \frac{1}{4}$ ,  $\epsilon < \frac{1}{2}(1-4\beta)$  and  $N$  is sufficiently large.

The contributions to (1.18) from  $T_\rho^{C,n}$  and  $T_\rho^{C,e}$  are handled similarly. To treat the contribution from  $T_\rho^{C,\infty}$ , we use the second estimate in (2.12) and

calculate (for (1.18))

$$\begin{aligned}
(2.16) \quad & \| |\xi''|(\sigma(\xi))^{-1}(\chi_2 \Delta \chi_1)^\wedge(\xi) \|_{L^2(T_\rho^{C,\infty})}^2 \\
& \leq C \iint_{|\xi| \geq 3|\rho|} \delta^{n-6} |\widehat{\chi}_2(\xi')|^2 |\widehat{\psi}_5(\delta \xi'')|^2 \frac{|\xi''|^2 d\xi' d\xi''}{|\xi|^4} \\
& \leq C \left( \int_{|\xi''| \leq |\rho|} \delta^{n-6} |\rho|^{-2\alpha-2} |\widehat{\psi}_5(\delta \xi'')|^2 |\xi''|^2 d\xi'' \right. \\
& \quad \left. + \int_{|\xi''| \geq |\rho|} \delta^{n-6} |\widehat{\psi}_5(\delta \chi'')|^2 |\xi''|^{-2\alpha} d\xi'' \right) \\
& = C \left( \delta^{-6} |\rho|^{-2\alpha-2} \int_0^{|\rho|\delta} |\widehat{\psi}_5(r)|^2 r^n \frac{dr}{r} \right. \\
& \quad \left. + \delta^{2\alpha-4} \int_{|\rho|\delta}^\infty |\widehat{\psi}_5(r)|^2 r^{n-2-2\alpha} \frac{dr}{r} \right) \\
& \leq C(\delta^{-6} |\rho|^{-2\alpha-2} + \delta^{2\alpha-4} (|\rho|\delta)^{-N}), \quad \forall N > 0,
\end{aligned}$$

which, for  $\delta = |\rho|^{-\beta}$  and  $N$  large is  $\leq C|\rho|^{-2\beta-2\epsilon}$  provided  $\beta < \frac{1}{4}$  and  $\epsilon < \alpha + 1 - 4\beta$ . A similar analysis holds for the  $T_\rho^{C,\infty}$  contribution to (1.19).

We now turn to controlling the  $q(x)u_0(x)$  terms in (1.17)–(1.19), as well as the contributions from the  $\Delta(\chi_0) \cdot \chi_1$  term in (2.1). Note that since  $q(x)$  is  $C^{n-1+\sigma}$  (for some  $\sigma > 0$ ),  $q(x)$  has an extension (see, e.g., [St70, Ch.6]) to a  $C^{n-1+\sigma}$  function of compact support on  $\mathbb{R}^n$ , which we also denote by  $q$ . The restriction of  $q$  to any  $\Pi \in M_{2,n}$  is still  $C^{n-1+\sigma}$ .

Let  $\{D_t : 0 < t < \infty\}$  be the one-parameter group of partial dilations on  $\mathcal{S}'(\mathbb{R}^{n*})$ ,

$$(D_t f)(\xi', \xi'') = t^{n-2} f(\xi', t\xi''),$$

which, for  $f, g \in L^1$ , satisfy  $\int_{\mathbb{R}^n} D_t f d\xi = \int_{\mathbb{R}^n} f d\xi$  and  $D_t(f * g) = D_t f * D_t g$ . Then

$$\begin{aligned}
(2.17) \quad \widehat{q} \widehat{u}_0(\xi) &= \widehat{q} * \widehat{u}_0(\xi) = D_\delta(D_{\delta^{-1}} \widehat{q}) * \delta^{-\frac{n-2}{2}} D_\delta(\widehat{\chi}_0(\xi') \widehat{\psi}_1(\xi'') e^{ix_0'' \cdot \xi''}) \\
&= D_\delta(D_{\delta^{-1}}(\widehat{q}) * \delta^{-\frac{n-2}{2}} \widehat{\chi}_0 \widehat{\psi}_1 e^{ix_0'' \cdot \xi''}).
\end{aligned}$$

Now, as  $\delta = |\rho|^{-\beta} \rightarrow 0$ ,  $D_{\delta^{-1}}(\widehat{q}) = \delta^{-(n-2)} \widehat{q}(\xi', \delta \xi'')$  converges weakly to the singular measure

$$(2.18) \quad Q(\xi') \otimes \delta(\xi'') = Q(\xi') d\xi',$$

where  $Q(\xi') = \int_{\mathbb{R}^{n-2}} \widehat{q}(\xi', \xi'') d\xi''$ ; note that  $q \in C^{n-1+\gamma}$  implies that the integral defining  $Q$  converges and  $Q$  satisfies  $(2.2)_{1+\gamma}$ . Letting  $F(\xi) = \widehat{\chi}_0(\xi') \widehat{\psi}_1(\xi'') e^{ix_0'' \cdot \xi''}$ , it follows from (2.17) that

$$\begin{aligned} (2.19) \widehat{qu}_0(\xi) &= D_\delta(D_{\delta^{-1}}(\widehat{q}) * \delta^{-\frac{n-2}{2}} F) \\ &= D_\delta((Qd\xi') * \delta^{-\frac{n-2}{2}} F) + D_\delta((D_{\delta^{-1}}\widehat{q} - Qd\xi') * \delta^{-\frac{n-2}{2}} F). \end{aligned}$$

If we define  $\widehat{\chi}_4(\xi') = Q *_{\mathbb{R}^2} \widehat{\chi}_0(\xi')$ , then  $\widehat{\chi}_4$  also satisfies condition  $(2.2)_{1+\gamma}$  (and thus  $(2.2)_{\alpha'}$  for  $0 < \alpha' < 1$ , so that (2.14) can be applied), and the first term in (2.19) is

$$(2.20) \quad D_\delta((Qd\xi') * \delta^{-\frac{n-2}{2}} F) = \widehat{\chi}_4(\xi') \delta^{\frac{n-2}{2}} \widehat{\psi}_1(\delta\xi'') e^{ix_0'' \cdot \xi''}.$$

Thus, the contributions to  $\|P_\delta(qu_0)\|_{L^2}$ ,  $\| |D''| \widetilde{G}_\rho(qu_0) \|_{L^2}$  and  $\|\widetilde{G}_\rho(qu_0)\|_{L^2}$  from the first term in (2.19) may be handled as the main  $\chi_2 \Delta \chi_1$  term was earlier, with the obvious absence of the factor  $\delta^{-2}$ . To control the contributions from the second term in (2.19), we use the elementary

**Lemma 5** *Let  $\varphi(x)$ ,  $f(x)$  be functions on  $\mathbb{R}^m$  such that  $\varphi(x)$ ,  $|x|\varphi(x)$ ,  $f(x)$  and  $|\nabla f(x)|$  are in  $L^1(\mathbb{R}^m)$ . Then,  $\forall \epsilon > 0$*

$$\begin{aligned} &\left| \left( \epsilon^{-m} \varphi\left(\frac{x}{\epsilon}\right) - \left( \int_{\mathbb{R}^m} \varphi dy \right) \delta(x) \right) * f(x) \right| \\ &\leq C_m (\|\varphi\|_{L^1} + \| |x|\varphi \|_{L^1}) \cdot (\|f\|_{L^\infty(B(0;|x|-1))} + \|\nabla f\|_{L^\infty(B(x;1))}) \cdot \epsilon. \end{aligned}$$

Applying this for  $\epsilon = \delta$ ,  $\xi' \in \mathbb{R}^2$  fixed, and using  $F \in \mathcal{S}$ ,  $|\widehat{q}(\xi)| \leq C(1 + |\xi|)^{-(n-1+\gamma)}$ , we find that,  $\forall N > 0$

$$(2.21) \quad |(D_{\delta^{-1}}(\widehat{q}) - Qd\xi') * F(\xi)| \leq C_N (1 + |\xi'|)^{-\gamma} (1 + |\xi''|)^{-N} \delta.$$

Hence, the second term in (2.19) is  $\leq C_N \delta^{\frac{n}{2}} (1 + |\xi'|)^{-\gamma} (1 + |\delta\xi''|)^{-N}$  and this allows the contributions to (1.17)–(1.19) to be dealt with as the  $\chi_2 \Delta_{x''} \chi_1$  term was before.

Finally, we need to establish the estimates (1.17–1.19) for the  $4|\rho| \overline{\partial} \chi_0$  term in (2.1); thus, we need to show

$$(2.22) \quad \|P_\rho(\overline{\partial} \chi_0 \cdot \chi_1)\|_{L^2} \leq C|\rho|^{-1-\epsilon},$$

$$(2.23) \quad \| |D''| \widetilde{G}_\rho(\overline{\partial} \chi_0 \cdot \chi_1)\|_{L^2} \leq C|\rho|^{-1-\beta-\epsilon}, \text{ and}$$

$$(2.24) \quad \|\widetilde{G}_\rho(\overline{\partial} \chi_0 \cdot \chi_1)\|_{L^2} \leq C|\rho|^{-1-2\beta-\epsilon},$$



for some  $\epsilon > 0$ . Using the fact that  $\widehat{\bar{\partial}\chi_0}(\xi')$  is rapidly decreasing and vanishes to first order at  $\xi' = 0$ , we may replace (2.6) with

$$\begin{aligned}
\|\widehat{\bar{\partial}\chi_0\chi_1}\|_{L^2(T_\rho^s)}^2 &\simeq \int_0^{\frac{|\rho|}{2}} \int_{B^2\left((0, \frac{r^2}{2|\rho|}); |\rho|^{-\frac{1}{2}-\epsilon_0}\right)} |\widehat{\bar{\partial}\chi_0}(\xi')|^2 d\xi' \delta^{n-2} |\widehat{\psi_1}(\delta r)|^2 r^{n-3} dr \\
&\leq c_N \left( \int_0^{\sqrt{2}|\rho|^{\frac{1-2\epsilon_0}{4}}} |\rho|^{-2-4\epsilon_0} \delta^{n-2} |\widehat{\psi_1}(\delta r)|^2 r^{n-2} \frac{dr}{r} \right. \\
&\quad + \int_{\sqrt{2}|\rho|^{\frac{1-2\epsilon_0}{4}}}^{\sqrt{2}|\rho|^{\frac{1}{2}}} \left(\frac{r^2}{2|\rho|}\right)^2 |\rho|^{-1-2\epsilon_0} \delta^{n-2} |\widehat{\psi_1}(\delta r)|^2 r^{n-2} \frac{dr}{r} \\
&\quad + \left. \int_{\sqrt{2}|\rho|^{\frac{1}{2}}}^{\frac{|\rho|}{2}} \left(\frac{r^2}{2|\rho|}\right)^{-N} |\rho|^{-1-2\epsilon_0} \delta^{n-2} |\widehat{\psi_1}(\delta r)|^2 r^{n-2} \frac{dr}{r} \right) \\
&\leq c_N \left( |\rho|^{-2-4\epsilon_0} \int_0^{\sqrt{2}|\rho|^{\frac{1-2\epsilon_0}{4}} \delta} |\widehat{\psi_1}|^2 r^{n-2} \frac{dr}{r} \right. \\
&\quad + |\rho|^{-3-2\epsilon_0} \delta^{-4} \int_{\sqrt{2}|\rho|^{\frac{1-2\epsilon_0}{4}} \delta}^{\sqrt{2}|\rho|^{\frac{1}{2}}} |\widehat{\psi_1}|^2 r^{n+2} \frac{dr}{r} \\
&\quad + |\rho|^{-1-2\epsilon_0+N} \delta^{2N} \int_{\sqrt{2}|\rho|^{\frac{1}{2}} \delta}^{\frac{|\rho|}{2} \delta} |\widehat{\psi_1}|^2 r^{n-2-2N} \frac{dr}{r} \Big) \\
&\leq c_N \left( (|\rho|^{-2-4\epsilon_0} + |\rho|^{-3-2\epsilon_0+4\beta} (|\rho|^{\frac{1-2\epsilon_0}{4}-\beta})^{-N'} \right. \\
&\quad \left. + |\rho|^{-1-2\epsilon_0+N-2N\beta-N'(\frac{1}{2}-\beta)}) \right)
\end{aligned} \tag{2.25}$$

for any  $N, N' \geq 0$ . As before, the contributions from  $T_\rho^n$  and  $T_\rho^e$  are handled similarly. Since  $\epsilon_0 < \frac{1}{2} - 2\beta$ , if  $N'$  is chosen large enough this yields (2.23) with  $\epsilon \leq 2\epsilon_0$ , which is weaker than the previously imposed  $\epsilon < \frac{1}{2}(1 - 4\beta)$ .

The desired estimates (2.23), (2.24) are even easier and hold for any  $\beta < \frac{1}{2}$ . The contribution to (2.24) from  $T_\rho^{C,s}$  is controlled as in (2.13), but with the factor  $\delta^{n-2}$  and with the  $\widehat{\chi_2}$  in the integrand replaced by  $\widehat{\bar{\partial}\chi_0}$ ; this is then dominated in the same manner as below (2.14). The  $T_\rho^{C,s}$  contribution to (2.25) is estimated as in (2.15), but with the absence of the  $\delta^{-4}$ . All other contributions are dealt with similarly.

This concludes the proof of Thm.4 for the case of potentials in the Hölder class  $C^{n-1+\sigma}(\overline{\Omega})$ ,  $\sigma > 0$ . The restrictions on  $\beta$  and  $\epsilon$  that we have needed are that  $\beta < \frac{1}{4}$  and  $\epsilon < \frac{1}{2}(1 - 4\beta)$ .

### 3 Remarks

(i) The proof of Thm. 4 needs to be slightly modified if we assume that the potential  $q(x)$  belongs to the Sobolev space  $H^{\frac{n}{2}+\sigma}(\Omega)$  for some  $\sigma > 0$ . Since  $\partial\Omega$  is Lipschitz, such a  $q(x)$  can, by the Calderón extension theorem, be extended to be in  $H^{\frac{n}{2}+\sigma}(\mathbb{R}^n)$ . Again denoting the extension by  $q$ , one has by Cauchy-Schwarz

$$(3.1) \quad \int_{\mathbb{R}^2} \left( \int_{\mathbb{R}^{n-2}} (1 + |\xi''|) |\hat{q}(\xi', \xi'')| d\xi'' \right)^2 (1 + |\xi'|)^\sigma d\xi' \leq c(\|q\|_{\frac{n}{2}+\sigma})^2$$

Thus,  $Q$  as in (2.18) belongs to  $L^2(\mathbb{R}^2; (1 + |\xi'|)^\sigma d\xi')$ , so that  $\widehat{\chi}_4 = Q *_{\mathbb{R}^2} \widehat{\chi}_0 \in L^2(\mathbb{R}^2; (1 + |\xi'|)^\sigma d\xi') \cap L^\infty$ . Replacing the uniform decay estimate (2.2) $_\alpha$  with

$$(3.2)_\sigma \quad \widehat{\chi}_2 \in L^2(\mathbb{R}^2; (1 + |\xi'|)^\sigma d\xi')$$

will allow us to handle the first term in (2.19). Furthermore, if for  $\xi'$  fixed, we let  $\phi(\cdot) = \widehat{q}(\xi', \cdot)$  in Lemma 5, then  $\phi(\xi'')$  and  $|\xi''|\phi(\xi'')$  are in  $L^1(\mathbb{R}^{n-2})$  with norms (as functions of  $\xi'$ ) in  $L^2(\mathbb{R}^2; (1 + |\xi'|)^\sigma d\xi')$ , and so the second term in (2.19) is  $\leq c_N \widehat{\chi}_6(\xi') (1 + |\delta\xi''|)^{-N}$ ,  $\forall N$ , with  $\widehat{\chi}_6$  satisfying condition (3.2) $_\sigma$ . So, we are reduced to repeating the analysis of Section 2 with (2.2) $_\alpha$  replaced by (3.2) $_\sigma$ . The decay of  $\widehat{\chi}_2$  was used in only two places in the argument. In (2.14), under (3.2) $_\sigma$ , we have the same estimate except for the absence of  $|\xi'|^{-2\alpha}$ ; however, this loss is absorbed into terms rapidly decreasing in  $|\rho|^{\frac{1}{2}}\delta = |\rho|^{\frac{1}{2}-\beta}$  where (2.14) is used. On the other hand, in (2.16) we may estimate the inner integral by

$$(3.3) \quad \int_{|\xi'| \geq 2|\rho|} |\widehat{\chi}_2(\xi')|^2 \frac{d\xi'}{(|\xi'|^2 + |\xi''|^2)^2} \leq \int_{\mathbb{R}^2} |\widehat{\chi}_2|^2 \frac{d\xi'}{(1 + |\xi'|)^\sigma |\xi'|^4} \\ \leq c|\rho|^{-4-\sigma} \text{ if } |\xi''| \leq \rho$$

and

$$(3.4) \quad \int_{\mathbb{R}^2} |\widehat{\chi}_2(\xi')|^2 \frac{d\xi'}{(|\xi'|^2 + |\xi''|^2)^2} \leq c|\xi''|^{-4} \text{ if } |\xi'| \geq \rho,$$

so that

$$\begin{aligned}
(3.5) \quad & \| |\xi''|(\sigma(\xi))^{-1}(\chi_2 \Delta \chi_1)^\wedge(\xi) \|_{L^2(T_\rho^{C,\infty})}^2 \\
& \leq C \left( \int_{|\xi''| \leq |\rho|} \delta^{n-6} |\rho|^{-4-\sigma} |\widehat{\psi}_5(\delta \xi'')|^2 |\xi''|^2 d\xi'' + \int_{|\xi''| \geq |\rho|} \delta^{n-6} |\widehat{\psi}_5(\delta \chi'')|^2 |\xi''|^{-2} d\xi'' \right) \\
& = C \left( \delta^{-6} |\rho|^{-4-\sigma} \int_0^{|\rho|\delta} |\widehat{\psi}_5(r)|^2 r^n \frac{dr}{r} \right. \\
& \quad \left. + \delta^{-2} \int_{|\rho|\delta}^\infty |\widehat{\psi}_5(r)|^2 r^{n-4} \frac{dr}{r} \right) \\
& \leq C_N (\delta^{-6} |\rho|^{-4-\sigma} + \delta^{-2} (|\rho|\delta)^{-N}) \\
& = C_N \left( |\rho|^{6\beta-4-\sigma} + |\rho|^{2\beta} (|\rho|^{\beta-\frac{1}{2}})^N \right), \quad \forall N,
\end{aligned}$$

which is  $\leq c|\rho|^{-2\beta-\epsilon}$  for  $N$  sufficiently large, since  $\beta < \frac{1}{2}$ . The restrictions on  $\beta$  and  $\epsilon$  are as before.

(ii) The construction of the approximate solutions given by Thm. 4 may be generalized by taking  $\chi_0$  to be an arbitrary analytic function of  $z = x_1 + ix_2$ , defined on a domain  $\Pi \cap \Omega \subset \subset \Omega' \subset \Pi$ . Since  $\bar{\partial}\chi_0 = \Delta_{x'}\chi_0 \equiv 0$  on  $\Omega$ , the resulting  $u = u_0 + u_1$  is still an approximate solution in the sense of Thm. 4, except that (1.8) no longer applies. Thus, Thm. 1 can be strengthened to conclude that  $(q_1 - q_2)|_\Pi$  is orthogonal in  $L^2(\Pi \cap \Omega, d\lambda_\Pi)$  to the Bergman space  $A^2(\Pi \cap \Omega)$  of square-integrable holomorphic functions on  $\Pi \cap \Omega$ . Furthermore, by repeating the construction using  $\bar{\rho} = \frac{1}{\sqrt{2}}|\rho|(\omega_R - i\omega_I)$ , which induces the conjugate complex structure on  $\Pi$ , for which the  $\bar{\partial}$  operator equals the  $\partial$  operator induced by  $\rho$ , we obtain that  $(q_1 - q_2)|_\Pi$  is also orthogonal to the conjugate Bergman space  $\overline{A}^2(\Pi \cap \Omega)$  of anti-holomorphic functions. (The analogue of this in two dimensions was obtained in [SU87b].) It would be interesting to make further use of this information.

(iii) To obtain variants of Thm. 1 establishing smaller sets of uniqueness in  $\partial\Omega$ , it might be useful to use approximate solutions associated to different two-planes. For this, it seems necessary to construct approximate solutions with much thinner supports, i.e., to overcome the restriction  $\beta < \frac{1}{4}$  in Thm. 4. Such an improvement might also be useful in extending the results to  $q_j \in L^\infty$ .

## References

- [A] G. Alessandrini, Stable determination of conductivity by boundary measurements, *Appl. Anal.*, **27** (1988), 153–172.

- [C] A. P. Calderón, On an inverse boundary value problem, in *Seminar on Numerical Analysis and its Applications to Continuum Physics*, Rio de Janeiro, 1980.
- [F] L. D. Faddeev, Growing solutions of the Schrödinger equation, *Dokl. Akad. Nauk SSR*, **165** (1965), 514–517 (trans. *Sov. Phys. Dokl.* **10** (1966), 1033.)
- [G] P. Grisvard, *Elliptic Problems in Nonsmooth Domains*, Pitman, Boston, 1985.
- [H65] S. Helgason, The Radon transform on Euclidean spaces, compact two-point homogeneous spaces and Grassmann manifolds, *Acta Math.* **113** (1965), 153–180.
- [H80] S. Helgason, *The Radon Transform*, Birkhäuser, Boston, 1980.
- [Ha96] P. Hähner, A periodic Faddeev-type solution operator, *Jour. Diff. Eqns.* **128** (1996), 300–308.
- [Is] V. Isakov, *Inverse Problems for Partial Differential Equations*, Springer-Verlag, Berlin, 1998.
- [KV84] R. Kohn and M. Vogelius, Determining conductivity by boundary measurements, *Comm. Pure Appl. Math.* **37** (1984), 113–123.
- [KV85] R. Kohn and M. Vogelius, Determining Conductivity by Boundary Measurements II: Interior Results, *Comm. Pure Appl. Math.* **38** (1985), 643–667.
- [N88] A. Nachman, Reconstruction from boundary measurements, *Annals of Math.* **128** (1988), 531–576.
- [N96] A. Nachman, Global uniqueness for a two-dimensional inverse boundary value problem, *Annals of Math.* **143** (1996), 71–96.
- [St70] E. M. Stein, *Singular Integrals and Differentiability properties of Functions*, Princeton Univ. Press, Princeton, 1970.
- [SU86] J. Sylvester and G. Uhlmann, A uniqueness theorem for an inverse boundary value problem in electrical prospection, *Comm. Pure Appl. Math.* **39** (1986), 92–112.

- [SU87a] J. Sylvester and G. Uhlmann, A global uniqueness theorem for an inverse boundary value problem, *Annals of Math.* **125** (1987), 153–169.
- [SU87b] J. Sylvester and G. Uhlmann, Remarks on an inverse boundary value problem, in *Pseudo-differential operators : proceedings of a conference held in Oberwolfach, February 2-8, 1986*, H. O. Cordes, B. Gramsch and H. Widom., eds., *Lecture notes in mathematics* **1256**, Springer Verlag, Berlin, 1987.
- [SU88] J. Sylvester and G. Uhlmann, Inverse boundary value problems at the boundary-continuous dependence, *Comm. Pure Appl. Math.* **41** (1988), 197–221.